

Biotics Research Corporation

Technical Support

Products #3500 & 3501

Thyrostim™

Nutritional Support for the Thyroid

Thyroid Function

The thyroid gland is actually a collection of individual glands (follicles). Here, newly synthesized hormone is secreted into a central lumen prior to release into the bloodstream. In general, thyroid hormones refer to T3 (triiodothyronine) and T4 (thyroxine). Though T4 is the main product, T3 is 3 to 4 times more active. T4 (with 4 atoms of iodine) is converted to T3 (with 3 atoms of iodine) via peripheral tissues, especially the liver and lung.

Parafollicular thyroid cells secrete calcitonin, a hormone employed in calcium and phosphate homeostasis. Calcitonin inhibits bone breakdown and accelerates bone calcium and phosphate uptake. Blood calcium levels control the secretion of calcitonin by a pathway independent of the pituitary gland. Thyroid hormones regulate metabolism and energy balance, as well as growth, development and activity of the nervous system.¹ Thyroid hormones stimulate carbohydrate and fat breakdown, increase protein synthesis, and increase the basal metabolic rate.

Several factors, including low metabolic rate, falling blood pressure, and conditions that increase the need for energy such as a cold environment, hypoglycemia, pregnancy or high altitude, stimulate the secretion of thyroid hormones. Thyroid hormones regulate their production via feedback mechanisms on the hypothalamus/pituitary axis. Thyroxine completes a negative feedback loop by blocking the release of the trophic hormone thyrotropin releasing hormone (TRH). High blood levels of estrogens and androgens decrease TRH production and thus thyroid function. Aging generally decreases glandular processes, including thyroid function.

Thyroid Hormone Synthesis

An individual consumes up to 500 mcg of iodine per day. Fully, one third is absorbed by the thyroid, where the synthesis of thyroid hormone begins, and the remainder is excreted. Iodine is oxidized by the thyroid to organic iodine via iodoperoxidase, an enzyme requiring hydrogen peroxide. The addition of iodine occurs at the 2,5 positions of tyrosine residues of thyroglobulin. Two di-iodotyrosine residues are coupled to form a thyroxine precursor on thyroglobulin. Secretion of thyroid hormones is initiated by the lysosomal degradation of thyroglobulin to release T3 and T4.

In the bloodstream, T3 and T4 are carried by thyroxine-binding globulin, thyroxine binding pre-albumin and serum albumin. It is the concentration of free (unbound) hormone that is important. T4 is converted to T3 and is taken up by appropriate target cells. Thyroid hormone is transported to the nucleus where it activates transcription. The actual thyroid hormone receptor is a chromosomal protein locked into the nucleus. Hormone binding activates the receptor as a transcription factor, resulting in the synthesis of multiple enzymes affecting metabolism. Thyroid hormone receptors also occur in mitochondria, so that thyroid hormones regulate oxygen consumption and ATP production directly. Brown fat-induced reactions to hypothermia are triggered by thyroid hormones.

Abnormal Levels of Thyroid Hormones

Varying degrees of hypothyroid function are routinely detected by laboratory tests and other measures.² Hypothyroidism is characterized by obesity, cold, dry skin and fatigue, while hyperthyroidism has opposite effects. Goiter and thyroid hyperplasia are consequences of abnormally low dietary iodine.


Nutritional Support

Iodine. The common form of iodine in foods is iodide. This is the reduced form of iodine. In thyroid tissue, follicles normally concentrate iodide 40-fold greater than blood concentrations. At maximal activity, the thyroid can contain up to 300 fold greater concentration of iodide than blood levels. Cells oxidize iodide to organically bound iodine, which is then chemically combined with tyrosine. Kelp is a natural source of iodine.

Tyrosine. This amino acid is a protein building block. In particular, many of the tyrosine residues of thyroglobulin are iodinated. Each molecule of thyroid hormone contains the equivalent of two tyrosine molecules. The uptake of tyrosine decreases with age.³

Biotics Research Bovine Neonatal Pituitary/Hypothalamic Glandular

Cytozyme-PT/HPT™ represents neonatal glandular tissue **Cytozyme-PT/HPT™**, as in all of Biotics Research Corporation's Cytozyme glandulars, represents a tissue concentrate, including peptides, proteins, nucleic acids and other nutrient factors. Biotics Research Corporation selected neonatal



tissues for several reasons. Neonatal glands and tissues possess very high anabolic activity. Factors associated with rapid growth are more likely to be present in neonatal glands than in adult glands. Histologic examination of adult (2-5 years) bovine glands and neonatal newborn bovine glands dramatically illustrates the differences between the two due to aging and environmental exposure.^{5,10} These factors contribute to a loss of organ function, accumulation of lipofuscin and increased fat accumulation in many organs with aging. In addition, neonatal tissues have not been subjected to long exposure to pollutants and environmental stressors. For example, independent evaluation of common pesticides in neonatal glandulars has indicated that levels are below the limits of detection. In addition, all glandular supplements produced by Biotics Research Corporation are obtained from domestic, USDA approved animals.

Minerals

Selenium. This trace mineral in the form of selenocysteine is required by a family of antioxidant enzymes, the glutathione peroxidases. These enzymes reduce peroxidized fatty acids in membranes to safe byproducts. On the other hand, selenium is also required for iodothyronine deiodinase, the enzyme located in peripheral tissues that is required to convert T4 to T3. Selenium deficiency decreases the iodothyronine de-iodinase activity. High iodine intake, when selenium intake is low, can lead to thyroid damage because thyroid glutathione peroxidase activity is reduced.⁴

Magnesium. Thyroid hormone is intimately associated with regulation of energy production and mitochondrial function. Indeed, mitochondria possess thyroid hormone receptors. Enzyme utilization of ATP generated by mitochondria requires



complex formation with magnesium, generally in a 1:1 ratio. Magnesium is essential for protein synthesis, cell replication and activation of the sodium-potassium pump, as well as regulation of calcitonin and parathyroid hormone.

Copper. Copper is a cofactor for the antioxidant enzyme, superoxide dismutase, and it is implicated in thyroid gland metabolism. Animal studies suggest that copper deficiency decreased hepatic mono-deiodinase,⁵ and that condition is associated with a reduction in T4 levels. Hormone reduction could be due to decreased T4 synthesis or to impaired T4 release.⁶

Manganese. This trace element can be considered an antioxidant because it is a cofactor for mitochondrial superoxide dismutase. Manganese complexes with ATP in certain energy requiring steps associated with carbohydrate metabolism. Very low manganese intake in lab animals retarded growth and thyroid hormone metabolism.⁷

Rubidium. A variety of anecdotal reports and case studies suggest that rubidium promotes thyroid metabolism. The mechanism is unknown.

Tyrosinase. This is a copper-containing enzyme and belongs to the family of monophenol oxidases. Tyrosinase is implicated in tyrosine pathways.

Vitamin A. Vitamin A is required for the normal development and maintenance of epithelial tissues. Both Vitamin A and thyroid hormone regulate gene expression, depending upon the organ.⁸ Furthermore, vitamin A supports the expression of receptors for thyroid and other hormones.⁹

Product Information: Thyrostim™ is available in bottles of 90 and 180 tablets.

Product Adjuncts: GTA™

For more information, contact our Client Services Department or one of our Technical Consultants at

Biotics Research Canada
Toll Free: (800) 840-1676
Box 283 Keswick ON L4P 3E2
Email: orders@bioticscan.com



References

1. Surks MI, Fevert R. Drugs and Thyroid Function. *New Engl J Med* 1995; 333: 1688-1694.
2. Lindsay RS. Hypothyroidism. *Lancet* 1997; 346: 413417.
3. Chen TS et al. Intestinal transport during the life span of the mouse. *J. Gerontol* 1990 45: B129- B133.
4. Hotz CS et al. Dietary iodine and selenium interact to affect thyroid hormone metabolism of rats. *J Nutr* 1997; 127: 1214-1218.
5. Kralik A et al. Concentrations of thyroid hormone in serum and activity of hepatic 5-monodeiodinase in copper-deficient rats. *Z. Ernährung Swiss* 1996; 35: 288-291.
6. Allen DK et al. Function of pituitary-thyroid axis in copper-deficient rats. *J Nutr* 1982; 112: 20432046.
7. Eder K et al. The effect of manganese supply on thyroid hormone metabolism on the off span of manganese depleted dams. *Biol Trace Elem Res* 1996; 55: 137-145.
8. Shull JD et al. Cell-type specific interaction between retinoic acid and thyroid hormone in the regulation of expressing of the gene encoding ornithine amino transferase. *Endocrinology* 1995; 136: 2120 2126.
9. Coustaut M et al. The influence of dietary vitamin A in triiodothyronine, retinoic and glucocorticoid receptor in liver of hypothyroid rats. *Br J Nutr* 1996; 176: 295-306.
10. Christensen F. Glandulars: comparison of sources. *Am Chiropractor* 1985; September.

Supplement Facts

Serving Size: 2 Tablets		Servings Per Container: 45
	Amount Per Serving	% Daily Value
Vitamin A (as palmitate)	2,500 IU	50%
Iodine (from kelp)	150 mcg	100%
Magnesium (as magnesium aspartate, gluconate, glycinate)	100 mg	25%
Selenium (from vegetable culture †)	50 mcg	71%
Copper (as copper gluconate)	1 mg	50%
Manganese (as manganese gluconate)	2 mg	100%
Rubidium (from vegetable culture †)	25 mcg	*
Lamb Pituitary/Hypothalamus complex concentrate (from ovine)	40 mg	*
Lactobacillus acidophilus (DDS-1)	2 million	*
Tyrosine (as L-Tyrosine)	100 mg	*
Tyrosinase (from mushroom)	10 units	*
Superoxide Dismutase (from vegetable culture †)	20 mcg	*
Catalase (from vegetable culture †)	20 mcg	*

*Daily Value not established

Other ingredients: Stearic acid (vegetable source), modified cellulose gum, cellulose, magnesium stearate (vegetable source) and food glaze.

† Specially grown, biologically active vegetable culture containing **Phytochemically Bound Trace Elements™** and naturally associated phytochemicals including polyphenolic compounds with SOD and catalase, dehydrated at low temperature to preserve associated enzyme factors.

RECOMMENDATION: One (1) to two (2) tablets each day as a dietary supplement or as otherwise directed by a healthcare professional.

KEEP OUT OF REACH OF CHILDREN

Store in a cool, dry area.
Sealed with an imprinted safety seal for your protection.

NDC #55146-03500 Rev. 5/08